The Walking Drugged



Kirk Hughes RN, EMT, CSPI

Costs of Substance Abuse

Abuse of tobacco, alcohol, and illicit drugs is costly to our Nation, exacting more than \$700 billion annually in costs related to crime, lost work productivity and health care.**

	Health Care	Overall
Tobacco	\$130 billion	\$295 billion
Alcohol	\$25 billion	\$224 billion
Illicit Drugs	\$11 billion	\$193 billion

—50 Years of Progress. A Report of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2014.

- -Centers for Disease Control and Prevention. Excessive Drinking Costs U.S. \$223.5 Billion. Accessed March 9, 2015.
- -National Drug Intelligence Center. National Drug Threat Assessment.

Chanhassen Class of 2016

2013 Dylan Turcotte: Heroin


2014 Macalob Bartram: Heroin


2015 Alex Snyder: DTP? Synthetic

2014 Carver County, Davis Colley: Synthetic cannabinoid contributed



PRESCRIPTION/OVER-THE-COUNTER VS. ILLICIT DRUGS*












The National Institute on Drug Abuse is a component of the National Institutes of Health, U.S. Department of Health and Human Services. NIDA supports most of the world's research on the health aspects of drug abuse and addiction. Fact sheets on the health effects of drugs of abuse and information on NIDA research and other activities can be found at www.drugabuse.gov.


Scope of the Problem



OD Fatalities in MN

MDH Report
MN 2015
572 Died from OD
411 Died from MVC

Since 2000 ~ 5000 died from OD



The Smoking Gun

HEROIN VS GUN DEATHS IN 2015

Heroin Deaths - 12,989 Gun Deaths - 12,979


CDC; THE WASHINGTON POST

2014 Drug OD Deaths


2010 & 2011 Oxycodone was at the top

2012 Heroin surpassed Oxycodone

Source of Info is Death Certificates.




Opioid death rate for every 100,000 by state (2015) I APPALACHIA I NORTHEAST




The Face of Opioid OD


The typical victim of opioid overdose is young, white and male.



*Includes ages "not stated" Note: The CDC notes that a person can have multiple causes of death, therefore the total number of deaths per drug may exceed the actual number of dead persons in the selected population. Data may not equal 100 due to rounding. Ages 14 and younger not charted amounted to 0.3%



Rx Pain Reliever Sources



Substance Abuse and Mental Health Services Administration, *Results from the 2013 National Survey on Drug Use and Health: Summary of National Findings*, NSDUH Series H-48, HHS Publication No. (SMA) 14-4863. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2014.

Heroin

Bayer heroin bottle.


From 1898 to 1910
 heroin was
 marketed as a non addictive morphine
 substitute and
 cough medicine for
 children!



The Jump from RX Opioids to Heroin


Painkillers And The Heroin Market

A growing number of people are using heroin in recent years, in part because it can be cheaper and easier to find than opioid painkillers purchased on the black market. Most heroin users were first hooked on prescription opioids, which generated \$11 billion in 2010 for the pharmaceutical industry.





Global Heroin Source and Distribution*




Novel Synthetic Opioids

OPIOIDS/ Analgesics

FENTANYL ACCOUNTED FOR APPROXIMATELY 70% OF THE OPIOID IDENTIFICATIONS. OF THE 90 IDENTIFICATIONS, FENTANYL WAS

FOUND AS THE ONLY CONTROLLED SUBSTANCE IN 30% OF THE IDENTIFICATIONS AND WITH HEROIN IN 40% OF THE IDENTIFICATIONS. Furanyl fentanyl was the next most

prominent opioid identified, accounting for approximately 10% of the identifications. Furanyl fentanyl was the only active substance reported in 50% of these identifications.




OPIOIDS/ Analgesics


FENTANYL ACCOUNTED FOR 50% OF THE IDENTIFICATIONS. THE NEXT MOST PROMINENT FENTANYL-RELATED SUBSTANCE, FURANYL FENTANYL, ACCOUNTED FOR 24% OF THE IDENTIFICATIONS.


ACRYL FENTANYL AND P-FLUOROBUTYRYLFENTANYL WERE REPORTED FOR THE FIRST TIME IN THE FOURTH QUARTER OF 2016.

The relative percentage of fentanyl identifications decreased from between 65 and 70% in previous quarters to 50% due to the significant increase (300%) in furanyl fentanyl identifications during the fourth quarter.



Furranyl Fentanyl (FuF)






Butyrfentanyl

- Analog of Fentanyl
- ¼ the potency of Fentanyl
- Not directly scheduled in the US.
- Onset 15-30 minutes,
 Duration 90-180 minutes



Acetyl Fentanyl

- Analog of Fentanyl
- 5-15x more potent than Heroin
- 2013 Rhode Island = 14 deaths
- 2013 Pennsylvania = 50 deaths
- Not technically Schedule
 1 in the US
- Onset 7-10 minutes,
 Duration 3-4 hours



Carfentanil

Wildnil


- Develop by JanssenPharmaceuticals 1974
- Large animal anesthetic
- 10,000 more potent than morphine
- Used in gas form by Russians in 2002 Moscow Theater Chechen hostage crisis 40 Terrorists killed & 130 Hostages died





MN & Carfentanil

- 6 Deaths directly related to adulterated Heroin
- 10 more possible
- Though Potent, lethality has not been established



Urban Myths with Carfentanil

- Dermal absorption causing opioid effects
- Inhalation of small incidental amounts causing significant opioid effects



DEA Quote

Oxycodone with "Pinky"

Research Chemical U-47700 in Counterfeit Oxycondone Pills

In March 2016, law enforcement officers in Lorain County, Ohio, seized 500 pills that visually appeared to be oxycodone. The pills were blue and had "A 215" markings, consistent with 30 milligram oxycodone pills (see Figure 5). Laboratory analysis indicated that the pills did not contain oxycodone, but were instead the research chemical U-47700. U-47700 is an unscheduled synthetic opioid not studied for human use that has caused at least 17 overdoses and several deaths in the United States. Although counterfeit opioid pill traffickers currently use fentanyls, it is likely that other synthetic opioids will be utilized if fentanyls become inaccessible, or if market prefences shift to other subtances.

Figure 5: Counterfeit Oxycodone Pills Containing U-47700.

Source: Lorain County, Ohio, Sheriff's Office

Fentanyl Deaths Double from 2013-2014

2013: 1,905 (6 per Million)

2014: 4,200 (13 per Million)



CDC Reports Increase in Fentanyl adulterated Opiates

- 2013-2014
- 426% Increase of seized Drugs testing (+) for fentanyl
- Data from 27 states
- Increase in number of deaths from Fentanyl & Synthetic opioids 79%
- No increase in Rx of Fentanyl during this period



Why Fentanyl Laced Heroin?

- Longer more potent high
 - Heroin is more euphoric, but has short ½ life.
 - Fentanyl = longer ½ life and in combination prolongs high
- Cheaper
 - 1 kilo of Heroin ~ \$35K
 - 1 kilo of Fentanyl ~3.5K



Gray Death

- New Street Slang a
 Mixture of any of the
 following: Heroin,
 U47700, Fentanyl,
 Carfentanil
- Routes
 - Ingested
 - Injected
 - Smoked
 - Snorted





Withdrawal

Opiate withdrawal generally not life threatening


 Can induce agitation- Safety issue for agency that cannot titrate naloxone doses



Other Trends

- Loperamide HCI
- Urban indications
 - Withdrawal
 - Abuse
- Unadvertised adverse effects
 - QRS widening/QTcProlongation





Kratom (Mitragyna speciosa)

Naturally occurring. Plant is dried then smoked, chewed, or made into a Tea/Capsules

Effects dose dependent

Low Dose = Stimulant Medium/High Dose = Opiate-like effects

No reported deaths from using only Kratom, but when mixed with OTC cold & cough meds or benzos, deaths have bee reported

Withdrawal reported; No likely life threatening: Agitation, pain, insomnia, fever, diarrhea, tremors, sweating. Generally last 1-3 days

Briefly listed by DEA as "Schedule 1" in 2016, but withdrawn shortly after



Minnesota Treatment

Methadone

- 15 Clinics Statewide
- ~6000 Patients

Suboxone

- Dr can dispense after 8 hour course
- 128 Certified in MN
- Previously manage 100 Patients, now 275

Kinetics

- ½ life of Buprenorphine = 24-48 hr
- ½ life of Naloxone = 2-12 hours





Robotrippin

- Seeking DEX (nonaddictive opiate effects in high amts)
- Coricidin HBP cough and cold also has antihistamine = anticholinergic effects
- Patients act intoxicated



Toxidromes






Opioid

- CNS depression
- Respiratory depression
- Miosis
- Hypothermia

TX: Narcan

Narcan also be effective in Clonidine, Tramadol

HOW TO GIVE NASAL SPRAY NARCAN

















MJ Kinetics

Smoked



Ingested



P450 Cytochrome



11-hydroxy-THC

Kinetics

Onset: Seconds to Minutes

Peak Effects: 15-30

Kinetics

Onset: 30-90 minutes

Peak: 1-6 hours*

147% in Peds MJ Exposures






Figure 1. Annual number and rate of marijuana exposures among children younger than 6 years (National Poison Data System 2000-2013).



Colorado Experience

Marijuana and Traffic Fatalities

Percent of All Traffic Deaths That Were Marijuana Related*



*Percent of All Fatalities Where the Operators Tested Positive for Marijuana

SOURCE: National Highway Transportation Safety Administration, Fatality Analysis Reporting System (FARS), 2006-2013 and CDOT/RMHIDTA 2014. NOTE: Cannot confirm 100% reporting prior to 2012.

New MJ Products





New MJ Products





MJ & Hyperemesis

MJ use can cause hyperemesis

TX: Capsaicin cream 0.075% on abdomen





MJ Ingestion Case Study

Omaha, NE: Aug 2016

53 y.o. male's wife calls 911

Unknowingly ate 4 brownies made by son w/ MJ

Crawling on floor using profanities and calling the cat "Bitch"



Dabs/Wax





Shatter is smooth, clear and solid.

>80 percent THC.



Budder or Wax has a creamy consistency from being whipped

70% THC.

Honey Oil

Liquid

THC % Varies

Wax Manufacture Hazards

3/25/15 Stearns County: 2 men, 18 and 19, were charged with third-degree murder in Stearns County on allegations that a fire that started while making wax killed an 85-year-old woman

Advantages:

- Easy to hide
- High THC content
- Can be "Vaped"
- Low Smell



UPDATE

- -One plead guilty to aiding & abetting 3rd degree murder: 7 years in prison
- -Other plead guilty: 7 months in jail & 25 years probation

MJ labs causing Explosions Nationwide

9/27/16: Explosion in the Bronx killing a FDNY Battalion Chief.

- Propane/Butane powering CO2 generators for accelerated MJ plant growth
- Precipitating hash oil from MJ extractors using butane while heating the precipitant





Synthetic Cannabinoids

Street Slang: Spice, K2, Klimax, Scooby Snax, Joker

12% of High Schoolers have tried in the last year

Multiple ingredients: Active compound mixed and sprayed on dried, leafy material

400+ synthetic variants exist = no consistent toxidrome

Paranoia, hallucinations, increased physical activity = Rhabdo, Acidosis, Hyperthermia, Renal Failure, Seizures


Addiction/Withdrawal potential unknown at this time




SYNTHETIC CANNABINOIDS

THERE WERE 88 SYNTHETIC CANNABINOID IDENTIFICATIONS IN THE THIRD QUARTER OF CY 2016. 5F-UR-144, AB-CHMINACA, AND

FUB-AMB ACCOUNTED FOR APPROXIMATELY 53% OF THE SYNTHETIC CANNABINOID IDENTIFICATIONS DURING THE REPORTING PERIOD.



SYNTHETIC CANNABINOIDS




SC Morbidity/Mortality

Source:

NEJM 373;2 Jul 9,2015

- N=810
- Mar 2012 Jan 2015
- Deaths = 23 or 3%

16 different SC homologs identified in the study



Bath Salts

Synthetic Cathinone

Also known as:

- Plant food
- Fertilizer

Wellbutrin structurally related





Synthetic Cathinone

- MDMA/MAMA (Ecstasy)
- Bath Salts
 - Alpha PVP (Flakka)
 - Ethylone
 - MDPV
 - Mephedrone (Meow Meow)
 - Methylone
 - BZP (Party Pills)
 - Many more
- 2C's: 2C-b, 2C-e, 2C-l








3rd Qtr 2016

CATHINONES

DIBUTYLONE WAS THE MOST COMMONLY IDENTIFIED CATHINONE, ACCOUNTING FOR APPROXIMATELY 25% OF THE IDENTIFICATIONS.



Dea.emerging.trends@usdoj.gov

Bath Salt Effects

- Hyperthermia
- Chest Pain
- Hypertension
- Tachycardia
- Excited Delirium
- Violent Behavior
- Foaming at the Mouth
- Limb Twitching
- Suicidal Thoughts
- Delusions



Acid, LSD, or NBOMe?

Analog of 2C-i/2C-c = 25C NBOMe Street Names

- N-Bomb
- Smiles
- Acid
- -Mistaken for LSD/Acid (Blotter)
- -Very Potent (16x more than parent compound) 50-250 mcg are purported dosages





Cocaine

- > 90% cut with Levamisole
- Old Vet dewormer
- Causes as an unintended side effects leukopenia (Low White Blood cells)



PCP...Its Back

- Street Names: Wet, Sherm. Illie
- Low use in the last decade +
- Disconnects pain receptors from the brain
- Very unpredictable & dangerous patients







Stuffer or Packer





Stuffer/Packer Obs

Meth = 24 hours





Heroin = 6 hours



Cocaine = 6 hours







Other Trends

Counterfeit Xanax & Oxycodone contain Fentanyl

- Xanax G3722 "Bars"
- Green Hulk





